

Einstein-Cartan-Dirac Theory In (1+2)-Dimensions

T. Dereli*

Department of Physics, Koç University
34450 Sarıyer, Istanbul, Turkey

N. Özdemir[†], Ö. Sert[‡]

Department of Physics Engineering, Istanbul Technical University
34469 Maslak, Istanbul, Turkey

Abstract

Einstein-Cartan theory is formulated in (1+2)-dimensions using the algebra of exterior differential forms. A Dirac spinor is coupled to gravity and the field equations are obtained by a variational principle. The space-time torsion is found to be given algebraically in terms of the Dirac field. Circularly symmetric, exact solutions that collapse to AdS_3 geometry in the absence of a Dirac spinor are found.

PACS numbers: 04.20.Jb, 04.60.Kz, 04.70.Bw

*tdereli@ku.edu.tr

†nozdemir@itu.edu.tr

‡sertoz@itu.edu.tr

1 Introduction

Field theories in (1+2)-dimensions often provide easy ways to check ideas that are difficult to prove in actual (1+3)-dimensions. However, it is not surprising to uncover other new ideas as well that are specific to (1+2)-dimensions. Topologically massive gravity [1] or BTZ black holes [2] are some of the best known examples to the latter. Other aspects and the literature may be found in Ref.[3]. In this paper we couple a Dirac spinor in (1+2)-dimensions to Einstein-Cartan gravity [4]. The field equations are derived by a variational principle. Then a family of circularly symmetric, rotating solutions that are asymptotically AdS_3 are found and discussed.

We specify the space-time geometry by a triplet (M, g, ∇) where M is a 3-dimensional differentiable manifold equipped with a metric tensor

$$g = \eta_{ab} e^a \otimes e^b \quad (1)$$

of signature $(-++)$. $\{e^a\}$ is an orthonormal co-frame dual to the frame vectors $\{X_a\}$, that is $e^a(X_b) = \delta_b^a$. A metric compatible connection ∇ can be specified in terms of connection 1-forms $\{\omega^a_b\}$ satisfying $\omega_{ba} = -\omega_{ab}$. Then the Cartan structure equations

$$de^a + \omega^a_b \wedge e^b = T^a, \quad (2)$$

$$d\omega^a_b + \omega^a_c \wedge \omega^c_b = R^a_b \quad (3)$$

yield the space-time torsion 2-forms $\{T^a\}$ and curvature 2-forms $\{R^a_b\}$, respectively. Here d denotes the exterior derivative and $\wedge$ the wedge product. We fix the orientation of space-time by choosing the volume 3-form $*1 = e^0 \wedge e^1 \wedge e^2$ where $*$ is the Hodge star map. It is possible to decompose the connection 1-forms in a unique way as

$$\omega^a_b = \hat{\omega}^a_b + K^a_b \quad (4)$$

where $\{\hat{\omega}^a_b\}$ are the zero-torsion Levi-Civita connection 1-forms satisfying

$$de^a + \hat{\omega}^a_b \wedge e^b = 0 \quad (5)$$

and $\{K^a_b\}$ are the contortion 1-forms satisfying

$$K^a_b \wedge e^b = T^a. \quad (6)$$

The curvature 2-forms are also decomposed in a similar way:

$$R^a_b = \hat{R}^a_b + \hat{D}K^a_b + K^a_c \wedge K^c_b \quad (7)$$

with

$$\hat{D}K^a_b = dK^a_b + \hat{\omega}^a_c \wedge K^c_b - \hat{\omega}^c_b \wedge K^a_c.$$

The field equations of Einstein-Cartan theory of gravity [4] are obtained by varying the action

$$I = \int_M (\mathcal{L}_{EC} + \mathcal{L}_M) \quad (8)$$

where the Einstein-Cartan Lagrangian density 3-form

$$\mathcal{L}_{EC} = -\frac{1}{2\kappa} R_{ab} \wedge *(e^a \wedge e^b) + \lambda * 1, \quad (9)$$

with the gravitational constant κ and the cosmological constant λ , and the matter Lagrangian density 3-form $\mathcal{L}_M$. We write the infinitesimal variations (up to a closed form) as

$$\dot{\mathcal{L}} = \dot{e}^a \wedge \left(-\frac{1}{2\kappa} R^{bc} * e_{abc} + \lambda * e_a + \tau_a \right) + \frac{1}{2} \dot{\omega}^{ab} \wedge \left(-\frac{1}{\kappa} * e_{abc} T^c + \Sigma_{ab} \right) \quad (10)$$

where the variations of the matter Lagrangian yield the stress-energy 2-forms

$$\tau_a = \frac{\partial \mathcal{L}_M}{\partial e^a} = T_{ab} * e^b \quad (11)$$

and the angular momentum 2-forms

$$\Sigma_{ab} = \frac{\partial \mathcal{L}_M}{\partial \omega^{ab}} = S_{ab,c} * e^c. \quad (12)$$

Therefore the Einstein-Cartan field equations are given either as

$$\frac{1}{2} R^{bc} * e_{abc} - \kappa \lambda * e_a = \kappa \tau_a, \quad (13)$$

$$* e_{abc} T^c = \kappa \Sigma_{ab} \quad (14)$$

or equivalently as

$$R_{ab} = \kappa \lambda (e_a \wedge e_b) + \kappa * e_{abc} \tau^c, \quad (15)$$

$$T_a = \frac{1}{2} \kappa * e_{abc} \Sigma^{bc}. \quad (16)$$

2 Einstein-Cartan-Dirac Field Equations

Let us consider a Dirac spinor field

$$\psi = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \quad (17)$$

and the conjugate spinor field

$$\bar{\psi} = \psi^\dagger \gamma_0 = (-\psi_2^* \ \psi_1^*) \quad (18)$$

where ψ_1 and ψ_2 are complex, odd Grassmann valued functions. We use a real (i.e. Majorana) realization of the gamma matrices $\{\gamma_a\}$ given explicitly as

$$\gamma_0 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \gamma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \gamma_2 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (19)$$

The exterior covariant derivative of the spinor fields are defined to be

$$\nabla\psi = d\psi + \frac{1}{2}\omega^{ab}\sigma_{ab}\psi \quad , \quad \overline{\nabla\psi} = d\bar{\psi} - \frac{1}{2}\omega^{ab}\bar{\psi}\sigma_{ab} \quad (20)$$

with

$$\sigma_{ab} = \frac{1}{4} [\gamma_a, \gamma_b] = \frac{1}{2} * e_{abc} \gamma^c. \quad (21)$$

We set $\gamma = \gamma_a e^a$.

Next let us introduce the (Hermitian) Dirac Lagrangian density 3-form

$$\mathcal{L}_D = \frac{i}{2} (\bar{\psi} * \gamma \wedge \nabla\psi - \overline{\nabla\psi} \wedge * \gamma \psi) + im\bar{\psi}\psi * 1. \quad (22)$$

Its infinitesimal variations are found to be (up to a closed form)

$$\begin{aligned} \dot{\mathcal{L}}_D = & \dot{e}^a \wedge \left\{ \frac{i}{2} * e^b_a \wedge (\bar{\psi} \gamma_b \nabla\psi + \overline{\nabla\psi} \gamma_b \psi) + im\bar{\psi}\psi * e_a \right\} \\ & + \frac{1}{2} \dot{\omega}^{ab} \wedge \left\{ \frac{i}{2} \bar{\psi} (*\gamma\sigma_{ab} + \sigma_{ab} * \gamma) \psi \right\} \\ & + i\dot{\bar{\psi}} \left\{ * \gamma \wedge \nabla\psi + \frac{1}{2} * e^a_b \wedge T^b \gamma_a \psi + m * \psi \right\} \\ & - i \left\{ \overline{\nabla\psi} \wedge * \gamma - \frac{1}{2} * e^a_b \wedge T^b \bar{\psi} \gamma_a - m * \bar{\psi} \right\} \dot{\psi}. \end{aligned} \quad (23)$$

From the above expression we identify the stress-energy 2-forms

$$\tau_a = \frac{i}{2} * e^b{}_a (\bar{\psi} \gamma_b \nabla \psi + \bar{\nabla} \bar{\psi} \gamma_b \psi) + im \bar{\psi} \psi * e_a \quad (24)$$

and the angular momentum 2-forms

$$\Sigma_{ab} = \frac{i}{2} (\bar{\psi} * \gamma \sigma_{ab} \psi + \bar{\psi} \sigma_{ab} * \gamma \psi). \quad (25)$$

Substituting these into the Einstein-Cartan equations (15) and 16) we obtain

$$\begin{aligned} R_{ab} = & \kappa \lambda e_a \wedge e_b + im \kappa \bar{\psi} \psi e_a \wedge e_b \\ & + i \frac{\kappa}{2} e_a \wedge (\bar{\psi} \gamma_b \nabla \psi) - i \frac{\kappa}{2} e_b \wedge (\bar{\psi} \gamma_a \nabla \psi) \\ & - i \frac{\kappa}{2} e_b \wedge (\bar{\nabla} \bar{\psi} \gamma_a \psi) + i \frac{\kappa}{2} e_a \wedge (\bar{\nabla} \bar{\psi} \gamma_b \psi), \end{aligned} \quad (26)$$

$$T_a = i \frac{\kappa}{2} \bar{\psi} \psi * e_a \quad (27)$$

and the Dirac equation

$$* \gamma \wedge \nabla \psi + m \psi * 1 = 0. \quad (28)$$

We note that with the torsion 2-forms (27), $*e^a{}_b \wedge T^b = 0$ identically and the Dirac equation simplifies to (28).

These field equations can be re-written in terms of the Levi-Civita connection only. To see this, we solve (27) for the contortion 1-forms

$$K_{ab} = -i \frac{\kappa}{4} \bar{\psi} \psi * (e_a \wedge e_b) \quad (29)$$

and substitute these into (26) which simplify to

$$\hat{R}_{ab} = \kappa \lambda e_a \wedge e_b + * e_{abc} \hat{\tau}^c - i \frac{\kappa}{4} d(\bar{\psi} \psi) \wedge * (e_a \wedge e_b) + \frac{3\kappa^2}{16} (\bar{\psi} \psi)^2 e_a \wedge e_b. \quad (30)$$

The Dirac equation (28) similarly simplifies to

$$* \gamma \wedge \bar{\nabla} \psi + m \psi * 1 - i \frac{3\kappa}{8} (\bar{\psi} \psi) \psi * 1 = 0 \quad (31)$$

It is interesting to note that the Einstein-Dirac equations (30) and (31) can be obtained from an action by zero-torsion constrained variations using the

method of Lagrange multipliers [5]. To this end, we consider a modified Dirac Lagrangian density 3-form

$$\mathcal{L}'_D = \frac{i}{2} \left(\bar{\psi} * \gamma \wedge \hat{\nabla} \psi - \overline{\hat{\nabla} \psi} \wedge * \gamma \psi \right) + im \bar{\psi} \psi * 1 + \frac{3\kappa^2}{16} (\bar{\psi} \psi)^2 * 1 \quad (32)$$

together with the constraint term

$$\mathcal{L}_{constraint} = (de^a + \omega^a_b \wedge e^b) \wedge \lambda_a \quad (33)$$

where $\{\lambda_a\}$ are the Lagrange multiplier 1-forms. The variation of the total action

$$I = \int_M (\mathcal{L}_{EC} + \mathcal{L}'_D + \mathcal{L}_{constraint})$$

with respect to the Lagrange multipliers imposes the constraint that the connection 1-forms are Levi-Civita. Then we solve the connection variation equations under this constraint for the Lagrange multiplier 1-forms as

$$\lambda_a = -\frac{i}{4} \kappa \bar{\psi} \psi e_a.$$

Substituting these in the remaining Einstein and Dirac equations give precisely (30) and (31).

3 A Circularly Symmetric Solution

We consider the metric

$$g = -f^2(r)dt^2 + h^2(r)dr^2 + r^2(d\phi + a(r)dt)^2 \quad (34)$$

in plane polar coordinates (t, r, ϕ) [6]. The following choice of orthonormal basis 1-forms

$$e^0 = f(r)dt \quad , \quad e^1 = h(r)dr \quad , \quad e^2 = r(d\phi + a(r)dt), \quad (35)$$

leads to the Levi-Civita connection 1-forms

$$\hat{\omega}^0_1 = \alpha e^0 - \frac{\beta}{2} e^2 \quad , \quad \hat{\omega}^0_2 = -\frac{\beta}{2} e^1 \quad , \quad \hat{\omega}^1_2 = -\gamma e^2 - \frac{\beta}{2} e^0 \quad (36)$$

where we defined

$$\alpha = \frac{f'}{fh} \quad , \quad \beta = \frac{ra'}{fh} \quad , \quad \gamma = \frac{1}{rh} \quad (37)$$

with ' denoting the derivative $\frac{d}{dr}$. On the other hand, assuming $i\frac{\kappa}{2}\bar{\psi}\psi = \tau(r)$, we calculate the contortion 1-forms

$$K^0_1 = \frac{\tau}{2}e^2, \quad K^0_2 = -\frac{\tau}{2}e^1, \quad K^1_2 = -\frac{\tau}{2}e^0. \quad (38)$$

As a first step towards a solution, we take a Dirac spinor that depends only on r and work out (28) in components as follows:

$$\begin{aligned} \psi'_1 + \frac{h}{2}(\alpha + \gamma)\psi_1 + \frac{h}{4}(\beta + 3\tau + 4m)\psi_2 &= 0, \\ \psi'_2 + \frac{h}{2}(\alpha + \gamma)\psi_2 + \frac{h}{4}(\beta + 3\tau + 4m)\psi_1 &= 0. \end{aligned} \quad (39)$$

We take the combinations $\psi_+ = \psi_1 + \psi_2$ and $\psi_- = \psi_1 - \psi_2$ and write a decoupled system of equations

$$\begin{aligned} \psi'_+ + (k_1 + k_2)\psi_+ &= 0 \\ \psi'_- + (k_1 - k_2)\psi_- &= 0 \end{aligned} \quad (40)$$

where we set

$$k_1 = \frac{h}{2}(\alpha + \gamma), \quad k_2 = \frac{h}{4}(\beta + 3\tau + 4m).$$

The formal solution to these equations are given by

$$\begin{aligned} \psi_1 &= e^{-\int^r k_1 dr} \left(\xi_1 e^{-\int^r k_2 dr} + \xi_2 e^{\int^r k_2 dr} \right) \\ \psi_2 &= e^{-\int^r k_1 dr} \left(\xi_1 e^{\int^r k_2 dr} - \xi_2 e^{-\int^r k_2 dr} \right) \end{aligned} \quad (41)$$

where ξ_1 and ξ_2 are complex, odd Grassmann valued constants. It can easily be verified

$$\tau(r) = i\kappa(\xi_2^* \xi_1 - \xi_1^* \xi_2) e^{-2\int^r k_1 dr}. \quad (42)$$

We next work out the Einstein field equations (15) which after simplifications reduce to the following system of coupled first order differential equations:

$$\begin{aligned} \frac{\beta'}{2h} + \beta\gamma &= -\frac{\tau'}{2h} - \tau\alpha \\ \frac{\gamma'}{h} + \frac{\beta^2}{4} + \gamma^2 + \kappa\lambda &= \frac{3\tau^2}{4} + \frac{\beta\tau}{2} \end{aligned}$$

$$\begin{aligned}
\frac{\alpha'}{h} - \frac{3\beta^2}{4} + \alpha^2 + \kappa\lambda &= \frac{3\tau^2}{4} - \frac{\beta\tau}{2} \\
\frac{\beta^2}{4} + \alpha\gamma + \kappa\lambda &= -\frac{3\tau^2}{4} - 2m\tau \\
\frac{\beta'}{2h} + \beta\gamma &= \frac{\tau'}{2h} + \tau\gamma.
\end{aligned} \tag{43}$$

At this point we choose

$$\kappa\lambda = -\frac{1}{l^2} < 0$$

and restrict our attention to those cases for which

$$\gamma = \alpha = \frac{1}{rh} \quad , \quad \tau = \beta = \frac{\beta_0}{r^2}.$$

We can then integrate (43) for the metric functions

$$f(r) = \frac{r}{l} \quad , \quad h(r) = \frac{l}{r\sqrt{1 - \frac{2m\beta_0 l^2}{r^2} - \frac{\beta_0^2 l^4}{r^4}}}, \tag{44}$$

and

$$a(r) = \frac{1}{2l} \arcsin\left(\frac{m}{\sqrt{m^2 + \frac{1}{l^2}}}\right) - \frac{1}{2l} \arcsin\left(\frac{m + \frac{\beta_0}{r^2}}{\sqrt{m^2 + \frac{1}{l^2}}}\right). \tag{45}$$

It now remains to integrate for the Dirac spinor and we find

$$\psi_{\pm} = \frac{1}{r} \left| r^2 - m\beta_0 l^2 + \sqrt{r^4 - 2m\beta_0 l^2 r^2 - \beta_0^2 l^4} \right|^{\pm \frac{ml}{2} \mp \frac{1}{2\beta_0} \arcsin\left(\frac{m + \frac{\beta_0}{r^2}}{\sqrt{m^2 + \frac{1}{l^2}}}\right)} e^{\pm \frac{1}{2\beta_0} \arcsin\left(\frac{m + \frac{\beta_0}{r^2}}{\sqrt{m^2 + \frac{1}{l^2}}}\right)} \xi_{\pm} \tag{46}$$

where $\xi_{\pm} = \xi_1 \pm \xi_2$.

In order to understand the physical meaning of this solution we write down the metric

$$g = -\frac{r^2}{l^2} dt^2 + \frac{l^2 dr^2}{r^2 \left(1 - \frac{2m\beta_0 l^2}{r^2} - \frac{\beta_0^2 l^2}{r^4}\right)} + r^2 (d\phi + a(r) dt)^2. \tag{47}$$

We see that in the absence of a Dirac spinor ($\beta_0 = 0$) the above metric collapses to the AdS_3 metric

$$g_0 = -\frac{r^2}{l^2} dt^2 + \frac{l^2}{r^2} dr^2 + r^2 d\phi^2. \tag{48}$$

Even when $\beta_0 \neq 0$, the metric $g \rightarrow g_0$ asymptotically as $r \rightarrow \infty$. We note a metric singularity at

$$r_c = \begin{cases} l\sqrt{m\beta_0 + \beta_0\sqrt{m^2 + \frac{1}{l^2}}} & , \quad \beta_0 > 0 \\ l\sqrt{|\beta_0|\sqrt{m^2 + \frac{1}{l^2}} - m|\beta_0|} & , \quad \beta_0 < 0 \end{cases} \quad (49)$$

We further calculate the curvature scalar

$$\mathcal{R} = -\frac{6}{l^2} + \frac{4m\beta_0}{r^2} \quad (50)$$

and the quadratic curvature invariant

$$R_{ab} \wedge *R^{ab} = \frac{6}{l^4} - \frac{8m\beta_0}{l^2 r^2} + \frac{\beta_0^2(8m^2 - \frac{4}{l^2})}{r^4} + \frac{16m\beta_0^2}{r^6} + \frac{8\beta_0^4}{r^8}. \quad (51)$$

The curvature invariants have an essential singularity at $r = 0$. We also check (See Ref.[6]) the quasi-local angular momentum

$$J(r) = \frac{r^3}{f(r)h(r)} \frac{da}{dr} = \beta_0, \quad (52)$$

the quasi-local energy

$$E(r) = \frac{1}{h_0(r)} - \frac{1}{h(r)} = \frac{r}{l} - \sqrt{\frac{r^2}{l^2} - 2m\beta_0 - \frac{\beta_0^2}{r^2}} \simeq \frac{\beta_0 m l}{r}, \quad (53)$$

and the quasi-local mass

$$\begin{aligned} M(r) &= 2f(r)E(r) = J(r)a(r) \\ &= 2\frac{l^2}{r^2} - 2\frac{l^2}{r^2}\sqrt{1 - \frac{2m\beta_0 l^2}{r^2} - \frac{\beta_0^2 l^2}{r^4}} \\ &\quad + \frac{1}{2l} \arcsin\left(\frac{m}{\sqrt{m^2 + \frac{1}{l^2}}}\right) - \frac{1}{2l} \arcsin\left(\frac{m + \frac{\beta_0}{r^2}}{\sqrt{m^2 + \frac{1}{l^2}}}\right) \\ &\simeq 2m\beta_0. \end{aligned} \quad (54)$$

4 Conclusion

We have formulated the Einstein-Cartan-Dirac theory in (1+2)-dimensions using the algebra of exterior differential forms. We coupled a Dirac spinor to Einstein-Cartan gravity and obtained the field equations by a variational principle. The space-time torsion is given algebraically in terms of the quadratic spinor invariant. We then looked for rotating, circularly symmetric solutions. We found a particular class of solutions that possesses an essential curvature singularity at the origin $r = 0$ that is hidden behind an event horizon at some finite distance $r = r_c$ away from the origin. The mass and the spin of this configuration can be identified. Thus the space-time geometry we found exhibits all the essential features of a black hole and we find it interesting that in the absence of the Dirac spinor field collapses to the regular AdS_3 geometry.

Acknowledgement

TD is supported in part by the Turkish Academy of Sciences (TÜBA).

References

- [1] S. Deser, R. Jackiw, S. Templeton, *Ann. Phys.* **140** (1982) 372
- [2] M. Banados, C. Teitelboim, J. Zanelli, *Phys. Rev. Lett.* **69** (1992) 1849
- [3] S. Carlip, **Quantum Gravity in (2+1)-Dimensions** (Cambridge University Press, 1998)
- [4] A. Trautman, *Bull. Acad. Polon. Scie.* **20** (1972) 185, 503, 895
- [5] T. Dereli, R. W. Tucker, *Class. Quant. Grav.* **5** (1988) 951
- [6] T. Dereli, Y. Obukhov, *Phys. Rev. D* **62** (2000) 0240013